

Ketamine

QUICK REFERENCE GUIDE

Indications	Dosing	PEAK EFFECT	PSYCHOLYTIC DOSE	PSYCHEDELIC DOSE
 Anxiety Bipolar disorder Depression Eating disorders OCD PTS Substance use disorders Suicidal ideation 	(70kg adult)	EFFECT	DOSE	DOSE
	IV INFUSION	1 min	35mg	70mg
	IM INJECTION	5 mins	33mg	75mg
	LOZENGE	20-30 mins	100mg	200mg
	PR SUPPOSITORY	20-45 mins	100mg	200mg
	PO LIQUID	1-2 hours	200mg	400mg
	RAPID DISSOLVE TABLET	15-30 mins	100mg	200mg
	INTRANASAL SPRAY (RACEMIC)	15 mins	75mg	150mg

Adverse Effects

- Changes in heart rate and blood pressure
- Increased salivation
- Nausea
- Vomiting
- Muscle spasms
- Headache
- Blurred vision
- Seizures
- Lower urinary tract symptoms
- Agitation
- Anxiety
- Increased suicidality

Subjective Effects

- Dissociation or a "trance-like" state
- Altered perceptions of time and space
- Changes in sensory perception
- Feelings of leaving one's body
- Emotionally intense visions
- Encounters with non-terrestrial beings
- Feelings of being reborn
- Vivid dreams or memories of past and future incarnations
- Ego dissolution and loss of identity
- Reliving one's life
- Deep feelings of peace and joy
- Feelings of interconnectedness with nature and others
- Feelings of cosmic unity



Absolute Contraindications

- Allergic reaction to ketamine
- Uncontrolled high blood pressure
- Acute angle glaucoma
- Uncontrolled thyroid disorder
- Pregnancy
- Suicide risk
- Current diagnosis of schizophrenia, schizoaffective disorder, major depressive disorder with psychotic features, psychotic disorder
- · Age 14 and under
- History of:
 - Ketamine abuse
 - · Stroke or brain aneurysm
 - Mania/bipolar

Relative Contraindications

- Active or unstable substance use disorder that is not an indication for treatment
- Dissociative identity disorder
- Current diagnosis of borderline personality disorder
- Active or chronic severe suicidality and self-injury
- Active breastfeeding
- Sleep apnea

History of:

- · Heart disease
- Psychosis
- Seizure disorders

Concomitant Medications

- Adderall (amphetamine / dextroamphetamine)
- Benadryl (diphenhydramine)
- Klonopin (clonazepam)
- Lyrica (pregabalin)
- Seroquel (quetiapine)
- Valium (diazepam)
- Xanax (alprazolam)